

Getting started with LORBRENA



Patient was compensated for her time.

Indication

LORBRENA is a prescription medicine that is used to treat adults with non-small cell lung cancer (NSCLC) that is caused by an abnormal anaplastic lymphoma kinase (ALK) gene, **and** that has spread to other parts of your body (metastatic).

Your healthcare provider will perform a test to make sure that LORBRENA is right for you. It is not known if LORBRENA is safe and effective in children.

Selected Safety Information

LORBRENA can cause serious side effects, including:

- **Liver problems due to interactions with other medicines.** Liver problems can be severe. It is important to know what medicines not to take during treatment with LORBRENA.

Please see Important Safety Information on pages 4 and 5. Please click for the [full Prescribing Information](#) and [Patient Information](#), or visit [LORBRENA.com](https://www.LORBRENA.com).

Getting started

If you and your doctor have decided that LORBRENA is right for you, this brochure can act as a guide to what you might expect while on treatment. Here, you will learn about some of the side effects you may experience while taking LORBRENA, as well as strategies that you can discuss with your healthcare team to help manage them.

Getting used to a new routine takes time. Working with your doctor, you can find strategies that support your treatment goals.

Contents

How to take LORBRENA	3
Important Safety Information	4-5
Your experience	6
When certain side effects happened in clinical trials and how common they were	8-9
Helping to manage certain side effects	10-14
Tips for everyday well-being	15
Side effect tracker	16-21
Resources and support	22-23

How to take LORBRENA® (lorlatinib)

Tell your healthcare provider about the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements.



Take LORBRENA exactly as your healthcare provider tells you to take it. Do not change your dose or stop taking LORBRENA unless your healthcare provider tells you to do so.



Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with LORBRENA if you develop side effects.



Swallow LORBRENA tablets whole. Do not chew, crush, or split LORBRENA tablets. Do not take LORBRENA tablets if they are broken, cracked, or not intact.



Take LORBRENA at approximately the same time each day.



You may take LORBRENA with or without food.



Avoid grapefruit or grapefruit juice while on treatment with LORBRENA.



If you miss a dose, take it as soon as you remember. However, if it is within 4 hours of the time to take your next dose, just take your next dose at your regular time.



If you vomit after taking a dose of LORBRENA, do not take an extra dose. Take your next dose at your regular time.

Important Safety Information

LORBRENA® (lorlatinib) can cause serious side effects, including:

- **Liver problems due to interactions with other medicines.** Liver problems can be severe. It is important to know what medicines not to take during treatment with LORBRENA.
- **Central nervous system (CNS) effects.** CNS effects can be severe. Tell your healthcare provider if you get any new or worsening symptoms of CNS effects, including: problems with thinking, such as forgetfulness or confusion, changes in mood, such as depression and thoughts about suicide or dying, psychotic effects, such as seeing or hearing things that are not real (hallucinations), seizures, changes in speech, and changes in sleep.
- **Increases in the cholesterol and triglycerides (lipid) levels in your blood.** Most people will get an increase in the lipid levels in their blood during treatment with LORBRENA.
 - If you get increases in the lipid levels in your blood, your healthcare provider may start you on a new medicine or increase your dose if you are already taking a medicine to lower the lipid levels in your blood.
 - Your healthcare provider will do blood tests to check the lipid levels in your blood before starting treatment, 1 and 2 months after starting treatment, and during treatment with LORBRENA.
- **Heart problems.** LORBRENA can cause very slow or abnormal heartbeats. Your healthcare provider will check your heart rhythm (electrocardiogram or EKG) before starting and during treatment with LORBRENA. In some people, these problems are severe, and you may need to get a pacemaker. Tell your healthcare provider right away if you feel dizzy or faint or have abnormal heartbeats.
- **Lung problems.** LORBRENA can cause severe or life-threatening swelling (inflammation) of the lungs during treatment that can lead to death. Symptoms may be like those from lung cancer. Tell your healthcare provider right away if you get any new or worsening symptoms of lung problems, including trouble breathing, shortness of breath, cough, or fever.
- **High blood pressure (hypertension).** Your healthcare provider will check your blood pressure before starting treatment, 2 weeks after starting treatment, and then at least every month during treatment with LORBRENA. Your healthcare provider may start or change your blood pressure medicine if you get high blood pressure. Tell your healthcare provider right away if you get signs or symptoms of high blood pressure, including headaches, dizziness, blurred vision, chest pain, or shortness of breath.
- **High blood sugar (hyperglycemia).** LORBRENA can cause new or worsening increases in your blood sugar levels. Your healthcare provider will do blood tests to check your blood sugar levels before starting and during treatment with LORBRENA. Your healthcare provider may start or change your blood sugar medicine if you get high blood sugar. Tell your healthcare provider right away if you get any signs and symptoms of high blood sugar, including: feeling very thirsty, needing to urinate more than usual, feeling very hungry, feeling sick to your stomach, feeling weak or tired, and feeling confused.

If you get serious side effects during treatment, your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with LORBRENA.

Before taking LORBRENA, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney problems
- have liver problems
- have had episodes of depression or seizures
- have high levels of cholesterol or triglycerides in your blood
- have problems with your heartbeat
- have lung or breathing problems
- have high blood pressure
- have diabetes or high blood sugar
- are pregnant or plan to become pregnant. LORBRENA can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with LORBRENA.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with LORBRENA.
- Use effective non-hormonal birth control during treatment and for at least 6 months after the final dose of LORBRENA.
- Birth control pills (oral contraceptives) and other hormonal forms of birth control may not be effective if used during treatment with LORBRENA. Talk to your healthcare provider about birth control choices that are right for you during this time.

Males with female partners who are able to become pregnant:

- Use effective birth control during treatment and for at least 3 months after the final dose of LORBRENA.

- are breastfeeding or plan to breastfeed. It is not known if LORBRENA passes into your breast milk. Do not breastfeed during treatment and for 7 days after the final dose of LORBRENA. Talk to your healthcare provider about the best way to feed your baby during this time.

Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements. Taking LORBRENA with certain other medicines may increase your risk of side effects and may affect the way LORBRENA or the other medicines work.

Know the medicines you take. Keep a list of them to show to your healthcare provider and pharmacist when you get a new medicine.

Do not take LORBRENA if you take certain other medicines called strong CYP3A inducers. Ask your healthcare provider for a list of these medicines if you are not sure.

The most common side effects of LORBRENA include:

- | | |
|---|--|
| • swelling in your arms, legs, hands, and feet (edema) | • pain in your joints |
| • numbness and tingling feeling in your joints or arms and legs (peripheral neuropathy) | • diarrhea |
| • weight gain | • changes in mood, such as depression and irritability |
| • problems with thinking, such as forgetfulness or confusion | • increased cholesterol and triglyceride levels in the blood |
| • tiredness (fatigue) | • cough |
| • difficulty breathing | |

LORBRENA may cause decreased fertility in males, which may affect your ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all of the possible side effects of LORBRENA.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Indication

LORBRENA is a prescription medicine that is used to treat adults with non-small cell lung cancer (NSCLC) that is caused by an abnormal anaplastic lymphoma kinase (ALK) gene, **and** that has spread to other parts of your body (metastatic).

Your healthcare provider will perform a test to make sure that LORBRENA is right for you.

It is not known if LORBRENA is safe and effective in children.

Your experience

Open communication and active partnership with your healthcare team is key. Let them know how you're feeling, including any side effects you may be experiencing.

Your healthcare team can work with you on possible ways to help manage side effects. Dose adjustments can be a part of treatment with LORBRENA.



The best advice I could give someone starting LORBRENA is to have open and honest conversations with your doctor about any side effects you experience, so you can work towards a plan to help manage them.

—Stephen, patient taking LORBRENA

Patient was compensated for his time.



LORBRENA[®] together

Taking LORBRENA? To sign up for resources and information via email or to request a starter kit, go to LORBRENA.com/sign-up



The most common side effects of LORBRENA[®] (lorlatinib) include:

- swelling in your arms, legs, hands, and feet (edema)
- numbness and tingling feeling in your joints or arms and legs (peripheral neuropathy)
- weight gain
- problems with thinking, such as forgetfulness or confusion
- tiredness (fatigue)
- difficulty breathing
- pain in your joints
- diarrhea
- changes in mood, such as depression and irritability
- increased cholesterol and triglyceride levels in the blood
- cough

These are not all of the possible side effects of LORBRENA.



If you have serious effects during treatment with LORBRENA, your doctor may change your dose, stop your treatment for a period of time, or completely stop treatment with LORBRENA.

Taking a lower dose of LORBRENA may help manage side effects.

Remember that everyone responds to treatment differently, so you may or may not experience some of the side effects described here.

Please see Important Safety Information on pages 4 and 5. Please click for the [full Prescribing Information](#) and [Patient Information](#), or visit LORBRENA.com.


When certain side effects happened in clinical trials and how common they were

How to understand the timeline


The side effects on this page are not all of the possible side effects of LORBRENA® (lorlatinib). The information here is based on certain side effects experienced in the LORBRENA clinical trials and does not include all of the side effects that were experienced by patients (see Important Safety Information on pages 4 and 5).

Likelihood of having side effect of any severity (based on clinical studies)

Low (0%-25%) Medium (26%-50%) High (51%-75%)






Likelihood of other common side effects



The values listed below represent the median (middle) time from the first dose of LORBRENA to when patients first reported the side effect. “Median” means about half of the patients reported it earlier and half reported it later than this time point.

You may or may not experience the side effects described below. If you do experience these side effects, you may not experience them in the same time frame or for the same amount of time.



For more information on the likelihood of side effects, please refer to the [full Prescribing Information](#).

Information about when certain side effects occurred is data from published medical journal articles.

Want to learn more about the clinical trials?
Visit [LORBRENA.com/clinical-study-results](https://lorbrena.com/clinical-study-results)



Please see Important Safety Information on pages 4 and 5. Please click for the [full Prescribing Information](#) and [Patient Information](#), or visit [LORBRENA.com](https://lorbrena.com).

LORBRENA®
LORLATINIB | 100 mg | 25 mg tablets

Managing certain side effects

See the information and tips below that you can discuss with your healthcare team to help manage certain side effects.



High cholesterol and/or triglycerides

What is it?

An increase in the levels of cholesterol and/or triglycerides in the blood

What can be done?

- Your oncologist or primary care physician will check your cholesterol and triglyceride levels by doing regular blood tests
- You may be prescribed a medicine to lower cholesterol, such as statins, and/or asked to consider lifestyle and diet changes. If needed, your doctor may increase the dose of your cholesterol-lowering medicine or prescribe a different medicine
- Let your doctor know about all the medicines and supplements you are taking. They will ensure everything works safely together



Changes in mood

What is it?

Feeling depressed or irritable

What can be done?

- If you or your care partner notice any change in your mood—even if it seems minor—talk to your doctor right away. Letting them know early can help protect your well-being
- Keep track of changes in mood by writing down how you feel: See the side effect tracker starting on page 16
- To manage these side effects, your doctor may pause treatment and/or reduce your LORBRENA® (lorlatinib) dose. In some cases, they may recommend seeing a mental health specialist
- In the clinical study, these side effects were often mild and, with proper management, reversible



Swelling (edema)

What is it?

Swelling of the arms, legs, hands, or feet

What can be done?

- Talk to your doctor if you notice any swelling. They can recommend strategies that work best for you
- Wear compression garments
- Elevate your legs and stay active to help reduce swelling
- Your doctor may tell you to limit salt intake
- To manage these side effects, your doctor may pause treatment and/or reduce your LORBRENA dose, or prescribe a diuretic



Psychotic effects (hallucinations)

What is it?

Smelling, hearing, or seeing things that are not real

What can be done?

- If you or your care partner notice any episodes of hallucination, talk to your doctor right away. Letting them know early can help protect your well-being
- To manage these side effects, your doctor may pause treatment and/or reduce your LORBRENA dose. In some cases, they may recommend seeing a mental health specialist
- In the clinical study, these side effects were often mild and, with proper management, reversible

Managing certain side effects (cont'd)



Tingling and numbness

What is it?

Pain, numbness, tingling, or muscle weakness

What can be done?

- Let your doctor know if you feel numbness, tingling, or discomfort
- A night splint can improve carpal tunnel syndrome
- Your doctor may recommend working with a physical therapist for pain or numbness
- Your doctor may tell you to limit salt intake
- To manage these side effects, your doctor may pause treatment and/or reduce your LORBRENA® (lorlatinib) dose, or prescribe vitamin B1 or B6
- In the clinical study, these side effects were typically reversible following a dose reduction



Changes in speech

What is it?

Trouble finding the right words or slurred speech

What can be done?

- Report any speech changes to your doctor as soon as you or your care partner notice them—even if they seem mild
- To manage these side effects, your doctor may pause treatment and/or reduce your LORBRENA dose
- In the clinical study, these side effects were often mild and, with proper management, reversible



Weight gain

What is it?

Increase in your body weight

Important Information:

- Treatment may increase your appetite, but there are also other factors that may cause you to gain weight
- Edema (swelling), which is extra fluid in the body, may play a role in weight gain

What can be done?

- Preventing weight gain is a helpful management strategy
- Talk with your doctor about nutrition, exercise, and other steps that have been proven to help manage weight gain and maintain your well-being
- Your doctor may check your weight regularly and suggest therapy management strategies that work for you

Understanding potential weight gain with LORBRENA

Scan the QR code or click [here](#) to learn about this possible side effect and working with your doctor to help address it.



Managing certain side effects (cont'd)



Problems with thinking

What is it?

Forgetfulness or confusion

What can be done?

- If you or your care partner notice any changes in memory or ability to think—even if they seem minor—talk to your doctor right away. Letting them know early can help protect your well-being
- Keep a diary or calendar with key information to remember: See the side effect tracker starting on page 16
- Create a system of keeping objects in the same place
- Your doctor may recommend working with a mental health specialist to treat certain behaviors
- To manage these side effects, your doctor may pause treatment and/or reduce your LORBRENA® (lorlatinib) dose
- In the clinical study, these side effects were often mild and, with proper management, reversible

Things to consider

Tips for everyday well-being

Set healthy goals

When you're living with ALK+ mNSCLC, healthy habits can help make a difference. Remember to always work with your doctor when making diet and lifestyle changes. Here are a few simple changes to think about.



Eat a balanced diet. What you eat may affect how you feel. If you're interested in eating healthier, talk to your doctor or a nutritionist.



Stay active. Light exercise and walking are activities you could ask your doctor about.



Rest. When you feel tired, listen to your body. Take a few moments to lie down or take a nap.



Join a support group. You're not alone. Explore support groups in your area or online. See page 22 for QR codes to advocacy organizations.



Keep track of your questions and how you feel. Write them down in a journal or notepad and take it with you to appointments to discuss with your healthcare provider.



Use the side effect tracker on page 16 to start.

Side effect tracker

Having an open conversation with your healthcare team and care partners can allow them to better help you during your treatment—they are your partners in helping you address side effects that may occur.

Below is a reminder of certain side effects you may experience. Use the side effect tracker on the next few pages to help your care team best understand how you are feeling by taking note of these or any other side effects.



High cholesterol and/or triglycerides



Changes in mood



Swelling (edema)



Psychotic effects (hallucinations)



Tingling and numbness



Changes in speech



Weight gain



Problems with thinking

Directions

To use the side effect tracker, write the date and each side effect you experience, and circle the number for how severe it is on a scale of 1 to 5.

- 1 means the side effect is minor, doesn't happen often, and doesn't last long. These side effects do not impact your daily routines
- 5 means the side effect is more severe and lasts a long time. These side effects impact your daily routines

Remember to make note of anything else you may be experiencing on that day.

Date	Name of side effect	Severity (circle one)				
		1	2	3	4	5

Notes:

Date	Name of side effect	Severity (circle one)				
		1	2	3	4	5

Notes:

Date	Name of side effect	Severity (circle one)				
		1	2	3	4	5

Notes:

Side effect tracker (cont'd)

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Side effect tracker (cont'd)

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Date	Name of side effect	Severity (circle one)
		1 2 3 4 5

Notes:

Resources and support

Below are a few helpful resources that provide tools, information, or support. Simply click the title of each resource or scan each code with your smartphone.



[Advocacy partners](#)

Discover organizations that provide helpful resources, like connection to support communities and information on the latest research.



[Patient stories](#)

See what people taking LORBRENA® (lorlatinib) are sharing about their diagnosis, their treatment journey, and what they're up to now.



[LORBRENA Together](#)

Request a starter kit and sign up for an email program to help you navigate your treatment journey.



[Additional helpful resources](#)

Find tools and resources to help you and your loved ones feel informed, supported, and ready for what's ahead.



[This Is Living With Cancer™](#) offers personalized support and resources designed to fit your needs. Sign up to access disease-specific articles, educational materials, cancer planning information, wellness content, and more.*



[For care partners](#)

Your support can make a real difference. Find tips and resources that can help you feel more prepared and connected.

*The free resources offered through **[This Is Living With Cancer™](#)** and **[LivingWith®](#)** are available to anyone living with cancer and their loved ones, and are not specific to LORBRENA.

Looking for support? We've got you covered

Pfizer Oncology Together™ understands that getting your prescribed Pfizer Oncology medicine is at the top of your to-do list. That's why our dedicated team offers you available resources, personalized support, and connections to financial assistance, if needed, to help you work through the challenges of access and reimbursement.

From finding financial assistance options to addressing coverage concerns, Pfizer Oncology Together is here to support you every step of the way.



Financial assistance

Eligible, commercially insured patients may pay as little as **\$0 per month** for their Pfizer Oncology treatment. **Limits, terms, and conditions apply.*** Patients may receive up to \$10,600 per product in savings annually.

The co-pay savings program provides assistance with out-of-pocket deductible, co-pay, or coinsurance costs.

*Patients are not eligible to use this card if they are enrolled in a state or federally funded insurance program, including but not limited to Medicare, Medicaid, TRICARE, Veterans Affairs health care, a state prescription drug assistance program, or the Government Health Insurance Plan available in Puerto Rico. For full Terms and Conditions, please visit PfizerOncologyTogether.com/terms.

Pfizer Oncology together™

Turn to Pfizer Oncology Together for financial assistance resources and personalized support.




Call **1-877-744-5675**
(Monday–Friday 8 AM–8 PM ET)

Visit **PfizerOncologyTogether.com**

Please see Important Safety Information on pages 4 and 5. Please click for the [full Prescribing Information](#) and [Patient Information](#), or visit LORBRENA.com.

LORBRENA®
LORLATINIB | 100 mg | 25 mg
tablets



For more information about
LORBRENA® (lorlatinib),
what to expect, support,
and resources, visit
[LORBRENA.com](https://lorbrena.com).

Patient was compensated for her time.

Please see Important Safety Information
on pages 4 and 5. Please click for the
[full Prescribing Information](#) and
[Patient Information](#), or visit [LORBRENA.com](https://lorbrena.com).

© 2026 Pfizer Inc. All rights reserved. July 2026 PP-LOR-USA-1217
LORBRENA® is a registered trademark of Pfizer Inc.


LORBRENA®
LORLATINIB | 100 mg
tablets

